MASCC International Cancer-Related Fatigue Guidelines



Strategies to Implement Exercise as a Treatment Modality to Manage Cancer-Related Fatigue

Stephen R Samuel, MPT, PhD
Assistant Professor, Dept. Of Physiotherapy,
Kasturba Medical College,
Manipal Academy of Higher Education, Mangalore
India



Conflicts of Interest

None to declare





Fatigue Guidelines-Exercise Group


Karen Mustian, PhD

Luke Peppone, PhD

Stephen Samuel, PhD







Summary of Studies

- Total Number of Studies Screened N = 128
- After Screening 91 studies
- Level of Evidence & Pedro Score was determined for each study
 - Level I N = 1
 - Level II N = 1
 - Level III N = 47
 - Level IV N = 42





Level I Exercise Study

One Level I Study M. Schmitz et al 2014

- Exercise may help overall and physical CRF
- Resistance with weight machines
- 60-80% of 1 RM
- 2 days /week
- 60 minutes
- 12 weeks
- Pedro Score: 8





Level II Exercise Study

One Level II Study / Winters-Stone 2012

- Null study regarding fatigue
- Anaerobic Resistance Exercise <u>vs</u> stretching
- Resistance with dumbells, barbells and weighted vests
- Moderate-Vigorous intensity
- 3/days week; two supervised; one at home
- 60 minutes
- 12 months
- Pedro Score: 8





Level III Exercise Studies

47 Level III Studies

- 25/47 had significant results
- Yoga, Aerobic , Resistance Training, Aerobic
 - + Resistance Training
- Avg. Pedro Score: 8





Positive Studies (Level III)- 25

- Length of intervention: 4 -12 Weeks
- Duration: 18 90 minutes
- Aerobic-9
- Combo(Aerobic + Resistance) 7
- Yoga 5
- Resistance Training 4



Level IV Exercise Studies

41 Level IV Studies





Summary

- Exercise thus as an intervention is <u>reasonable</u> to be <u>suggested</u> as an intervention to alleviate cancer related fatigue.
- The following exercise interventions have Level
 3 evidence & hence reasonable to suggest
 - 1. Aerobic exercise training
 - 2. Combination of Aerobic & Resistance training
 - 3. Yoga
 - 4. Resistance Training
 - 5. Walking





Strategies

- Awareness amongst oncologists/ physicians & patients
- Exercise & Psychological interventions found to be more effective (Mustian et al 2017)
- Exercise program accessibility remains a challenge in implementing the guidelines
- Delivered by certified & trained PT/ OT/ Exercise scientists
- Insurance





Pathway Model

- Referral, screening, medical clearance, and programming for people
- Mina et al 2018





- Exercise screening and risk stratification
- Appropriate instruction in exercise by a QEP
- Provision of resources to support exercise engagement (for example, manuals, log books; Web sites or apps; and basic exercise equipment such as resistance bands, stability ball)

 Routine follow-ups with the QEP, scheduled to ensure appropriate adjustment of the exercise prescription to accommodate changes in capacity, goals, or health status



- Mode & Dose
- Aerobic
- Aerobic + Resistance
- Yoga
- Intensity: Moderate
- Duration: 20-60 minutes







Questions ???

